

Circulation of Marginal and Semi-Enclosed Seas (Sea of Japan and Related Process Studies)

Christopher N.K. Mooers

Ocean Prediction Experimental Laboratory (OPEL)

Division of Applied Marine Physics

Rosenstiel School of Marine and Atmospheric Science

University of Miami

4600 Rickenbacker Causeway

Miami, FL 33149-1098

phone: (305) 361-4088 fax: (305) 361-4701 email: cmooers@rsmas.miami.edu

Grant Number: N000140110261

<http://www.rsmas.miami.edu/divs/amp.html>

LONG-TERM GOAL

My long-term goal is to understand the circulation dynamics of marginal and semi-enclosed seas through numerical simulation. Understanding the weather-driven transient flows (especially in coastal regions), mesoscale variability, ventilation, seasonal and interannual variability, and flow interactions with the basin topography is part of this goal.

OBJECTIVES

I wish to determine the “necessary and sufficient” conditions for usefully accurate numerical simulations as a prelude to the implementation of nowcast/forecast systems, which requires attention to model evaluation using observations as well as other models. For example, we need to establish the space-time resolution and amplitude accuracy requirements for atmospheric forcing of marginal and semi-enclosed seas. Given the difficulty of determining, in particular, open (lateral) boundary conditions, it is anticipated that data assimilation will be required.

APPROACH

We are using the Princeton Ocean Model (POM) as implemented on a mesoscale-admitting grid (ca. 10 km resolution) and with 26 sigma levels (and with relatively high, logarithmic resolution in the surface and bottom boundary layers) for the Japan (East) Sea (JES). It is driven with surface wind stress, heat flux, and moisture flux, and with specified throughflow from the Korea/Tsushima Strait to Tsugaru and Soya Straits. The model output is compared to available data, including CREAMS I (1993 thru 1996) current meter data (from Prof. Masaki Takematsu, Kyushu U.) over the Japan Basin and now CREAMS II (1999 thru 2001) data. Using synoptic forcing, simulation datasets for the CREAMS II field experimental period are generated and used for model evaluations. Upon completion of the model evaluations, data will be assimilated through the CREAMS II period to form an analysis that can be used for diagnostic studies of the JES dynamics and the analysis of alternative observing system strategies.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2002		2. REPORT TYPE		3. DATES COVERED 00-00-2002 to 00-00-2002	
4. TITLE AND SUBTITLE Circulation of Marginal and Semi-Enclosed Seas (Sea of Japan and Related Process Studies)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Ocean Prediction Experimental Laboratory (OPEL),,Division of Applied Marine Physics,Rosenstiel School of Marine,and Atmospheric Science,University of Miami,Miami,,FL, 33149				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

In my research group (OPEL), Ms. (now Dr.) HeeSook Kang defended her dissertation on simulations of wintertime ventilation and subduction in the JES; Dr. Inkweon Bang is simulating the CREAMS II period with synoptic (NOGAPS) forcing, evaluating model-data comparisons, and designing data-assimilative analyses; and Dr. Francisco Sandoval conducted model-data comparisons using CREAMS II simulations from Dr. Bang and CREAMS II observations, beginning with shipborne CTD data (from Prof. Lynne Talley, SIO) and Palace Float CTD and flow data (from Prof. Steve Riser, Univ. of Washington).

WORK COMPLETED

Synoptic atmospheric (NOGAPS) and monthly throughflow (NRL) forcing drove several simulation cases covering the CREAMS II period. The parametric dependence of these cases was a factor in model-data comparisons and corresponding sensitivity studies.

Three simulation JES-POM cases and GDEMS (NRL) climatology (as a baseline case) were compared to the Lynne Talley summer and winter CTD-grids and the ca. 21/2-year Steve Riser Palace Float CTD profiles and flows.

Dr. HeeSook Kang joined the University of Southern Mississippi (SSC Campus) as a NRL post doc.

Two mss were submitted to the CREAMS II special issue of Deep-Sea Research and are listed under PUBLICATIONS below.

Dialogue continued, with several other American, Japanese, Korean, and Russian JES modelers and observationalists, for coordination of model-model and model-data comparisons with CREAMS I and II observations.

RESULTS

Substantial agreement exists between the numerical simulations with JES-POM and the shipborne and Palace Float CTD data for the upper ocean temperature and salinity fields, more so in winter than summer and for the winter/summer difference fields. The main discrepancies are associated with the strength of the seasonal thermocline and halocline. Both the simulations and Palace Float flow data at 800 m define a large cyclonic recirculation gyre over the Japan Basin; however, the simulated speeds of the mean flow were about twice those observed. The patterns of speed variance were similar in the simulations and the observations; however, the simulated variance was generally smaller than observed.

The impacts of surface atmospheric forcing of different time-space scales on the simulation of water mass formation and spreading of formed water were investigated by quantifying water mass subduction/formation/transformation rates for the JES. It was found that, without synoptic atmospheric forcing, the diagnostics using JES-POM may underestimate, by a factor of two, buoyancy loss at the surface, and, hence, water mass formation, as well as mixing and spreading of the formed water mass.

IMPACT/APPLICATIONS

The model-data comparisons presented for CREAMS I and CREAMS II timeframes demonstrate that simulations with synoptic forcing can be rationally compared with available observations; i.e., the

simulations are in sensible ranges and available observations are useful for these purposes. Furthermore, the point has been reached where (1) useful feedback is being provided to the simulation efforts from the comparisons with observations, (2) the confidence in the verisimilitude of the simulations is sufficient to diagnose aspects of the wintertime ventilation and subduction, and (3) useful assessments of observing systems are in prospect.

TRANSITIONS

These results will be presented in Seoul at the mid-October 2002, NSF-sponsored, bilateral USA/Korea workshop on the JES and nowcast/forecast systems, which will form the basis for follow-on collaborative research in the topic area.

RELATED PROJECTS

OPEL is evolving a nowcast/forecast system for Prince William Sound, which is sponsored by the Oil Spill Recovery Institute (OSRI) of Cordova, Alaska, and another for the East Florida Shelf/Straits of Florida as part of the SEA-COOS Program.

REFERENCES

None

PUBLICATIONS

(2001) (with S. L. Vaughan and S. M. Gay III) Physical Processes Influencing the Pelagic Ecosystem of Prince William Sound. *Fisheries Oceanography*, 10 (Supplement 1), pp. 58-80.

(2001) (with J. Wang, M. Jin, V. Patrick, J. Allen, S. Vaughan, D. Eslinger, and T. Cooney) Numerical Simulations of the Seasonal Ocean Circulation Patterns and Thermohaline Structures of Prince William Sound, Alaska. *Fisheries Oceanography*, 10 (Supplement 1), pp. 132-148.

(2001) (with K.A. Korotenko Prediction of the Dispersal of Oil Transport in the Caspian Sea Resulting from a Continuous Release. *Spill Sci. Technol. Bull.*, and R.M. Mamedov) 6 (5/6), pp.323-329.

(2001) (with J. Wang, M. Jin, V. Patrick, J. Allen, S. Vaughan, D. Eslinger, and T. Cooney) Numerical Simulations of the Seasonal Ocean Circulation Patterns and Thermohaline Structures of Prince William Sound, Alaska. *Fisheries Oceanography*, 10 (Supplement 1), pp. 132-148.

(2002) (with L. Gao, W. D. Wilson, W. E. Johns, K. D. Leaman, H. E. Hurlburt, and T. Townsend) Initial Concepts for IAS-GOOS. In: Operational Oceanography : Implementation at the European and Regional Scales (Ed. N. C. Flemming, S. Vallergera, N. Pinardi, H. W. A. Behrens, G. Manella, D. Prandle, J. H. Stel). Elsevier, New York, pp. 379-390.

(2002) Global Geophysical Monitoring Systems. In: Volume 1, The earth system: physical and chemical dimensions of global environmental change (Eds. M.C. MacCracken and J.S. Perry), Encyclopedia of Global Environmental Change, John Wiley & Sons, pp. 534-539.

(2002) (with I. Bang) The Influence of Several Factors Controlling the Interactions between Prince William Sound, Alaska and the Northern Gulf of Alaska. *J. Phys. Ocean.* (in press).

(2002) (with I. Bang and S.L. Vaughan) Initial Steps Toward Validation of a Seasonal Cycle Simulation for Prince William Sound, Alaska. Circulation (Flow and Mass) Fields. *Con. Shelf Res.* (submitted).

(2002) Coastal Oceanography, *UNESCO Encyclopedia of Life Support Systems* (submitted).

(2002) (with H. Kang) Diagnoses of Simulated Water Mass Subduction/Formation/Transformation in the Japan/East Sea (JES). *Deep-Sea Res.* (submitted).

(2002) (with I. Bang and F. Sandoval) Synoptic Comparisons of Numerical Simulations with CREAMS II Observations of JES Circulation (Flow and Mass Fields). *Deep-Sea Res.* (submitted).

(2002) (with C. Andrade and D. Barton) Evidence for an Eastward Undercurrent along the South American Caribbean Coast) *J. Geophys. Res.* (submitted).

PATENTS

None